## **Supplementary Information for:**

# The impact of diet on the composition and relative

## abundance of rumen microbes in goat

4 Kaizhen Liu, Qin Xu, Lizhi Wang\*, Jiwen Wang, Wei Guo, Meili Zhou

5

1

3

Institute of Animal Nutrition, Sichuan Agricultural University, Ya'an, 625014, China

7

8

6

Table S1a. The dietary composition and nutrient contents for goats fed at a complete

#### 9 feed (CF) diet (dry matter basis<sup>1)</sup>)

Ingredients	Content (%)	Nutrient levels of diet (%) <sup>3)</sup>	
Alfalfa meal	35	Crude protein	11.9
Chinese wildrye	35	Calcium	0.6
Corn	11	Phosphorus	0.3
Soybean meal	5	Neutral Detergent Fiber	38.9
Wheat bran	13	Acid Detergent Fiber	15.4
Premix <sup>2)</sup>	0.5	Ethanol Extract (%)	5.2
NaCl	0.5		
Total	100		

<sup>1)</sup> Data were computed based on dry matter, the same below.

13 3) All nutrients are measured values.

14

15

16

17

<sup>2)</sup> premix provides: Fe(as ferrous sulfate) 30 ppm; Cu(as copper sulfate) 10 ppm; Zn(as

zinc sulfate) 50ppm;Mn(as manganese sulfate) 60ppm;VA2937IU;VD343IU;VE30IU.

## Table S1b. The dietary composition and nutrient contents for goats fed at all forage

### 19 (AF) diet (dry matter basis).

Ingredients	Content (%)	Nutrient levels of diet (%)	
Chinese wildrye	100	Crude Protein	9.0
Total	100	Calcium	0.95
		Phosphorus	0.3
		Neutral Detergent Fiber	64.3
		Acid Detergent Fiber	44.1
		Ethanol Extract	1.7

- Fig. S1 The number of OTUs detected in goats fed at a completed feed (CF) or all forage
- 34 (AF) diet.

